

The WiggleZ Dark Energy Survey

ABSTRACT

The accelerating expansion of the universe, attributed to “dark energy”, has no accepted theoretical explanation. The origin of this phenomenon unambiguously implicates new physics via a novel form of matter exerting negative pressure or an alteration to Einstein’s general relativity. These profound consequences have inspired a new generation of cosmological surveys that will measure the influence of dark energy using various techniques. One of the forerunners is the WiggleZ Survey at the Anglo-Australian Telescope, a new large-scale high-redshift galaxy survey that is now 50% complete and scheduled to finish in 2010. The WiggleZ project is aiming to map the cosmic expansion history using delicate features in the galaxy clustering pattern imprinted 13.7 billion years ago. In this article we outline the survey design and context, and predict the science highlights.

Chris Blake and the WiggleZ team highlight the design and targets of this innovative survey.



A perplexing picture of cosmology has emerged over the last decade. Several different observations concur that the universe has entered a startling phase of accelerating expansion propelled by a mysterious repulsive component of gravity known as “dark energy”. This component apparently constitutes about 70% of the current energy density of the universe. Intriguingly, the cosmic acceleration began at roughly the same time as the beginnings of life on our planet. The study of dark energy has a fascinating history, as reviewed recently in these pages by Calder and Lahav (2008).

There is currently no theoretical explanation for the existence or magnitude of dark energy. The accelerating cosmos therefore provides an outstanding opportunity for cosmologists to challenge fundamental physics, with two possible outcomes. First, the universe could



1 (a): The robot positioner placing optical fibres on the field plate of the AAOmega spectrograph. (b): The Anglo-Australian Telescope, situated near the Warrumbungle National Park. (Both B Norris)

2: The distribution of galaxies currently observed in the three WiggleZ Survey fields located near the Northern Galactic Pole. The observer is situated at the origin of the co-ordinate system. The radial distance of each galaxy from the origin indicates the observed redshift, and the polar angle indicates the galaxy right ascension. The faint pattern of galaxy clustering is visible in each field.




3: Behaviour of the scale factor of the universe, a , for a cosmological constant model with matter density $\Omega_m=0.3$ and cosmological constant density $\Omega_\Lambda=0.7$. (Top): Variation with redshift of the scale factor. (Middle): Variation with redshift of the time derivative of the scale factor, \dot{a} , which represents the “velocity” of the cosmic expansion. The y-axis is in units of H_0 , the Hubble constant at $z=0$. (Bottom): Variation with redshift of the second time derivative of the scale factor, \ddot{a} , which represents the “acceleration” of the cosmic expansion. The y-axis is in units of H_0^2 . Notice that the transition from decelerating to accelerating expansion occurs at redshift $z \sim 0.7$, which lies near the centre of the target redshift range of the WiggleZ Survey, as illustrated in the middle plot.

be dominated by a new and unsuspected form of matter that exerts a negative pressure. The leading candidate here is the “cosmological constant”, the energy of the quantum vacuum. However, it is extraordinarily difficult for fundamental quantum theory to reproduce the observed amplitude of dark energy; typical predictions overestimate it by more than one hundred orders of magnitude. These problems have motivated alternative suggestions in which dark energy is codified as a “scalar field” that fills all space. Examples of other hypothesized scalar fields in physics include the mechanism that propelled cosmic inflation, and the Higgs field, which may endow particles with mass.

The second possible outcome is that Einstein’s vision of gravity, general relativity, is incorrect on large cosmological scales. In this case dark energy would describe the discrepancy between relativity and an unknown modified theory of gravity, for which a number of suggestions exist. Alternatively, perhaps our derivation of the equations of cosmology, in which we approximate the universe as homogeneous despite its extreme clumpiness, can induce the illusion of accelerating expansion (Wiltshire 2007).

In short, dark energy is a precious clue in the quest to understand fundamental theory. Unravelling its origin is likely to entail a revolution in our understanding of physics, string theory or quantum gravity. A rich landscape of cosmological surveys has grown in recent years to meet this goal, measuring the influence of dark energy via a variety of observational techniques. A combination of observations is in fact required to identify unequivocally the nature of dark energy. In particular, we must seek out discrepancies between the expansion history of the universe and the rate of growth of cosmic structure within that overall expansion. This comparison can distinguish between the different propositions of dark energy such as modified gravity and scalar fields.

4: The redshift range and cosmic volume mapped by the WiggleZ survey (red line) compared to various current (black) and future (green) galaxy redshift surveys: 2dFGRS, 2-deg Field Galaxy Redshift Survey; SDSS-DR6, Sloan Digital Sky Survey; BOSS, Baryon Oscillation Spectroscopic Survey to be undertaken by the Sloan Consortium; WFMOS, survey performed by the proposed Wide-Field Multi-Object Spectrograph, see Bassett *et al.* 2005.



Galaxy redshift surveys

The WiggleZ Dark Energy Survey at the Anglo-Australian Telescope is one of the earliest of this new generation of dark-energy focused surveys. It commenced in August 2006 and is scheduled to finish in July 2010. The observations are being performed by a small core team of 14 Australia-based astronomers. The goal of the project is to construct a new large-scale galaxy redshift survey spanning a deep and wide slab of the cosmos from moderate-redshift ($z \sim 0.25$) to high-redshift ($z \sim 1$) over a sky area of 1000 square degrees. Part of the current galaxy database is displayed in figure 2.

This unprecedented combination of redshift depth and survey area will allow us to probe the nature of dark energy with multiple methods, covering the redshift range where the transformation from decelerating to accelerating expansion is thought to take place (see figure 3). Our position in the parameter space of current and future galaxy redshift surveys is illustrated in figure 4. WiggleZ is much deeper in redshift than previous wide-area spectroscopic surveys such as the 2-degree Field Galaxy Redshift Survey (2dFGRS) and the Sloan Digital Sky Survey (SDSS). It is orders of magnitude wider in areal coverage than previous high-redshift surveys such as the Deep Extragalactic Evolutionary Probe (DEEP2).

The large-scale structure of the universe is one of the most fertile sources of cosmological information and is a long-standing pillar of cosmology's "standard model". Patterns of galaxy clustering, shaped by the competing forces of inward gravity and outward expansion, contain information about the constituents of the universe and about the physical processes by which cosmic structure condenses and amplifies from initial small fluctuations. In the early 1990s, combined analyses of galaxy clustering and the Cosmic Microwave Background (CMB) fluctuations provided the first strong

evidence for the currently accepted dark-energy dominated cosmological model (Efstathiou *et al.* 1990, Ostriker and Steinhardt 1995). The model received spectacular confirmation from later observations of faint supernovae as standard candles (Riess *et al.* 1998, Perlmutter *et al.* 1999).

The cosmological power of galaxy redshift surveys was boosted further by the realization that the pattern of galaxy clustering encoded a robust "standard ruler" that could be used to map out the cosmic expansion history in a manner analogous to supernova standard candles (Eisenstein *et al.* 1999, Blake and Glazebrook 2003, Seo and Eisenstein 2003).

The nature of this standard ruler is a small preference for pairs of galaxies to be separated by a co-moving distance of 150 Mpc. This favoured scale is an extraordinary echo of sound waves that propagated 13.7 billion years ago through the matter-radiation plasma before CMB last-scattering, less than 380 000 years after the Big Bang. These sound waves emanated from primordial dark matter haloes, which launched spherical wave-crests driven by radiation pressure from the compressed baryon-photon plasma. These sound waves travelled rapidly in the early universe, at 58% of the speed of light, covering a co-moving distance of 150 Mpc between the Big Bang and the last-scattering epoch. Both the initial dark matter halo and the spherical baryonic shell preferentially seed the later formation of galaxies, imprinting the standard ruler into the large-scale clustering pattern. Because the radiation is tightly coupled to the baryons, the signature of the acoustic waves is also imprinted in the microwave background photons (see figure 5).

A powerful aspect of this cosmological distance probe that differentiates it from supernova observations is that the standard ruler can be applied in both the tangential and radial directions, i.e. perpendicular and parallel to our

Baryon oscillations as a standard ruler

Galaxies cluster with a known preferred separation, ΔS_{BAO} , which is determined from the physics of the CMB. In our observations of the galaxy distribution this preferred length imprints itself as a preferred angular separation on the sky, $\Delta \theta_{\text{BAO}}$, and a preferred redshift separation, ΔZ_{BAO} . Measurements of $\Delta \theta_{\text{BAO}}$ and ΔZ_{BAO} at different redshifts can be used to deduce the cosmic distance-redshift relation, $D_A(z)$, and Hubble-parameter-redshift relations, $H(z)$:

$$D_A(z) = \Delta S_{\text{BAO}} / \Delta \theta_{\text{BAO}} \quad (1)$$

$$H(z) = c \Delta Z_{\text{BAO}} / \Delta S_{\text{BAO}} \quad (2)$$


The measurements of D_A and H may be linked to the underlying contents of the universe that determine the expansion history. For a geometrically flat universe containing dark energy with an equation of state w :

$$H(z) = H_0 \sqrt{[\Omega_m(1+z)^3 + (1-\Omega_m)(1+z)^{3(1+w)}]} \quad (3)$$

$$D_A(z) = \int_0^z [c/H(z')] dz' \quad (4)$$

where Ω_m is the current matter density, H_0 is Hubble's constant, and c is the speed of light. Thus measurements of $H(z)$ and $D_A(z)$ may be used to deduce the properties of dark energy.

line-of-sight (see box "Baryon oscillations as a standard ruler"). Tangentially we measure a preferred angular clustering scale of about 3° , which determines the cosmic distance in a similar manner to a supernova flux. Radially we extract a preferred redshift-space clustering scale, which specifies the cosmic expansion rate or Hubble parameter at high redshift. This expansion parameter depends directly on the underlying energy densities and is extremely



5: Baryon acoustic oscillations measured in the CMB radiation and galaxy distribution. (Left): The spectrum of temperature fluctuations in the CMB, as determined by the Wilkinson Microwave Anisotropy Probe (fig. 12 of Bennett *et al.* 2003 *ApJS* 148 1, Spergel *et al.* 2003). The “wiggles” are produced by acoustic waves in the baryon–photon plasma before recombination. The peak in the power spectrum at an angular scale of approx 1° corresponds to the angular size of the 150Mpc sound horizon projected at the distance of the CMB. **(Right):** The spectrum of density fluctuations in the distribution of luminous red galaxies in the Sloan Digital Sky Survey (fig. 3 of Eisenstein *et al.* 2005). The bump at separation $105h^{-1}\text{Mpc}=150\text{Mpc}$ corresponds to a preferred clustering scale and is the corresponding imprint of the acoustic waves in the galaxy distribution. The amplitude of this signal is much lower than that seen in the CMB radiation because the baryons, which carry the imprint of acoustic oscillations, are sub-dominant to cold dark matter.

difficult to measure with other methods.

The standard ruler scale of 150Mpc can be predicted very accurately from the densities of dark matter and baryons inferred from observations of the CMB fluctuations. The Wilkinson Microwave Anisotropy Probe (WMAP) has already measured the scale with an error of 1.3% (Komatsu *et al.* 2008), and this will be improved even further by the European Space Agency’s Planck satellite, which will soon be launched. The difficulty in applying the ruler lies in measuring the signal over vast cosmic distances. The accuracy with which a galaxy survey can measure the underlying spectrum of clustering fluctuations is limited ultimately by its volume, which determines the number of Fourier modes of a given scale that the survey can resolve. In Fourier space, the 150Mpc preferred scale corresponds to a series of harmonics (“wiggles” or “baryon oscillations”) in the galaxy power spectrum, by analogy with the set of frequency harmonics excited by an organ pipe of fixed length. It is interesting to note that current galaxy redshift surveys have mapped less than 0.01% of the volume information (or Fourier amplitudes) in the observable universe.

The baryon oscillation standard ruler has already been detected at low redshifts ($z=0.2$ and $z=0.35$) in the distribution of luminous red galaxies in the SDSS (Eisenstein *et al.* 2005, see figure 5). The aim of the WiggleZ Dark Energy Survey is to extend this delineation of the redshift–distance relation by adding further data points up to $z=1$. After the addition of the WiggleZ results, which equate to a roughly 2% measure of the cosmic distance scale at $z=0.7$, the accuracy with which baryon oscillation data

can probe the properties of dark energy will be similar to supernova data (see figure 6). This will provide a detailed cross-check of the two techniques, in which tensions could be indicative of systematic errors or new physics. Indeed, existing comparisons of the SDSS baryon oscillation measurements with supernova data are already showing hints of disagreements (Percival *et al.* 2007). The WiggleZ Survey will also allow us to measure the growth of structure and test modified gravity theories.

Designing the WiggleZ Survey


The key design decision for a galaxy redshift survey is how to select the spectroscopic targets. From a cosmological viewpoint, galaxies are simply tracer particles of the underlying spectrum of density fluctuations that encodes the baryon oscillation and growth information. Different classes of galaxy vary in the details of how they trace the density fluctuations. For example, red quiescent elliptical galaxies inhabit dense cluster environments more often than blue star-forming spiral galaxies. However, on large cosmological scales, these details of galaxy formation are unimportant and can be essentially reduced to a single number, a “galaxy bias factor” specific to each galaxy class. The decision of which galaxy to target can then be simplified to observational considerations such as exposure time required to obtain a successful redshift.

The WiggleZ Survey is obtaining redshifts for more than 200 000 luminous blue star-forming galaxies. The spectra of these galaxies, illustrated in figure 7, are dominated by patterns of strong atomic emission lines originating in the ionized nebular gas in star-forming regions (in

particular the rest-frame lines [OII] $\lambda=3727\text{\AA}$, H β $\lambda=4861\text{\AA}$ and [OIII] $\lambda=4959\text{\AA}$, 5007\AA). These emission lines provide a galaxy redshift in a relatively short exposure of 1 hour on the 3.9m Anglo-Australian Telescope, avoiding the need to detect the galaxy “continuum” light, which is about four magnitudes fainter than the spectroscopic targets observed by 2dFGRS or SDSS. The primary target database for the WiggleZ Survey is provided by the orbiting Galaxy Evolution Explorer (GALEX) satellite, a NASA Small Explorer Class mission, which is mapping the sky in ultraviolet (UV) light with λ between 1400\AA and 2800\AA , usually blocked by the Earth’s atmosphere. Star-forming galaxies produce prodigious amounts of UV emission from their population of hot, young stars. The WiggleZ and GALEX collaboration is delivering about 1000 orbits of new UV imaging data across our target fields. In order to produce a sharp angular position for follow-up spectroscopy, we match the UV catalogues with optical imaging from the SDSS and the Red Cluster Sequence (RCS2) database obtained at the Canada-France-Hawaii Telescope.

We obtain WiggleZ redshifts using a multi-object fibre spectrograph at the Anglo-Australian Telescope called AAOmega, which is one of the world’s most complex and successful astronomical instruments (Lewis *et al.* 2002, Saunders *et al.* 2004, Sharp *et al.* 2006; see figure 1). AAOmega can obtain up to 392 galaxy spectra simultaneously using optical fibres placed on the focal plane by a robot positioner. The large field-of-view of the instrument, a 2° diameter circle on the sky, provides an unrivalled mapping speed. The new dual-beam spectrographs

6 (Top): A simulation of the power spectrum of galaxy clustering that will be measured from the final WiggleZ Dark Energy Survey. The power spectrum has been divided by a smooth "reference" spectrum fit in order to illustrate the detection of the baryon oscillations in the clustering pattern. (Bottom): Simulated cosmological parameter measurements using baryon oscillations in the final WiggleZ Dark Energy Survey. This figure focuses on measurements of the matter density, Ω_m , and the equation of state of dark energy, w . We assume a fiducial cosmological constant model with $\Omega_m=0.27$ and $w=-1$. The yellow ellipse indicates the 68% confidence region for measurement of these parameters using the WiggleZ Survey baryon oscillations combined with the CMB "acoustic scale parameter" ℓ_A , which calibrates the baryon oscillation preferred scale. The orange band indicates the confidence region obtained from WMAP measurements of the CMB "shift parameter". The red region displays the confidence region for latest supernova measurements. When the WiggleZ survey is complete, the baryon oscillation data will measure the properties of dark energy with a similar precision to the supernova data, providing a detailed cross-check of the two techniques. The central confidence circles illustrate the dark energy measurements obtained by combining all the datasets.



were commissioned in February 2006 to replace the original spectrographs, which had been used for the 2dFGRS. Bench-mounted in the stable West Coude room at the AAT, the new spectrographs feature a wider simultaneous wavelength coverage, higher spectral resolution and improved sensitivity. The WiggleZ collaboration has also provided a new dichroic to extend the wavelength coverage of the low-resolution mode redwards to 9500 Å in order to detect the highly redshifted emission lines of our targets.

Completing the full WiggleZ Survey will require 160 clear nights of telescope time, spread over four years, mapping a total area of 1000 square degrees split into several fields. At the time of writing (July 2008) the WiggleZ Survey is roughly 50% complete, having expended 112 nights to gather 110 000 galaxy redshifts. The first major scientific results will be announced shortly. The survey is also committed to public data releases of spectra and redshifts, the first scheduled for late 2008.

Expected science highlights

The WiggleZ Survey will have a substantial impact beyond the measurement of baryon oscillations. Firstly, the survey will yield an accurate measurement of the shape of the galaxy clustering power spectrum on large scales. This shape depends on the relative abundance of baryons and dark matter in the early universe, and can be used in conjunction with the CMB to provide more accurate estimates of the composition of the universe. Of particular interest is that massive neutrinos act as hot dark matter in the early universe, free-streaming to smooth out patterns of structure. While particle physics neutrino oscillation experiments accurately determine the mass difference between neutrino species, cosmological observations can yield the absolute mass.

Secondly, the survey will test theories of gravity by mapping the growth of structure with redshift. This is achieved by extracting the coherent patterns of galaxy velocities that result

from the bulk flow of matter into clusters and superclusters. These velocities cause small distortions in the measured galaxy redshifts, which collectively imprint a characteristic signature in the derived clustering spectrum. The strength of this signature, in comparison to the overall clustering amplitude of the galaxies, allows us to track the growth rate of structure.

Thirdly, detailed analysis of the star-formation rates, environments, morphologies and luminosity functions of the WiggleZ dataset will provide information on galaxy evolution. These objects are typically forming stars at a rate of tens of solar masses per year, significantly faster than our own Milky Way. The highest-redshift galaxies in the sample have prodigious star-formation rates exceeding $100 M_{\odot} \text{yr}^{-1}$. Do these massive starbursts result from major galaxy mergers? Is this process suppressed or enhanced in dense cluster environments? Finally, the hundreds of thousands of high-redshift galaxy spectra contain statistically significant samples of rare and

interesting classes of object such as AGNs.

In parallel with our programme of observations, we are generating large-scale cosmological N -body simulations using the supercomputer at Swinburne University of Technology. These simulations, populated with star-forming galaxies, will be a crucial ingredient in our interpretation of the survey data.


Future prospects

The next decade is expected to be a golden age of cosmology with a series of impressive new surveys. WiggleZ will be succeeded by the Baryon Oscillation Spectroscopic Survey (BOSS), which should be executed by the upgraded SDSS telescope between 2009 and 2014. BOSS will target a vast volume of luminous red galaxies over 10 000 square degrees out to redshift $z=0.7$ and also plans to map higher-redshift structure along the lines-of-sight to high-redshift quasars. In the medium term, there is a currently unfunded proposal to build a powerful 5000-fibre Wide-Field Multi-Object Spectrograph for an 8 m telescope. At higher redshifts the Hobby-Eberly Telescope Dark Energy Experiment (HETDEX) plans to harvest spectra of Lyman- α emitting galaxies. Future radio telescopes will detect the faint glow of neutral hydrogen emission from high-redshift galaxies, enabling transformational radio redshift surveys.

In parallel, a new suite of deep and wide optical imaging surveys will emerge. The first Panstarrs telescope on Mauna Kea is already operating and another three are planned. The Dark Energy Survey project will be mapping 5000 square degrees of southern sky from the 4 m Blanco telescope in Chile. A dedicated 8 m wide-field Large Synoptic Survey Telescope (LSST) is on the horizon. These deep images will allow cosmologists to detect millions of new supernovae and achieve precise measurements of the faint signature of gravitational lensing. New space missions have also been proposed, including the Joint Dark Energy Mission (JDEM) in the United States and the Euclid mission in Europe.

This wealth of new data will pose an unprecedented challenge to the existing standard cosmological model. If the unexpected need for dark energy is interpreted as a serious failure of the model, then we could be on the brink of a paradigm shift that introduces a new cosmological framework. ●

The WiggleZ team: Chris Blake, Sarah Brough, Warrick Couch, Karl Glazebrook, Greg Poole (Swinburne University of Technology); Tamara Davis, Michael Drinkwater, Russell Jurek, Kevin Pimblett (University of Queensland); Matthew Colless, Rob Sharp (Anglo-Australian Observatory); Scott Croom (University of Sydney); Michael Pracy (Australian National University); David Woods (University of New South Wales); Barry Madore (OCIW); Chris Martin, Ted Wyder



7: Example spectra from the survey. (a) Galaxy with a high-quality redshift ($z=0.6628$) determined by multiple emission lines. (b) Galaxy with a high-quality redshift ($z=0.3048$) determined by multiple lines, at a lower redshift than (a). (c) Galaxy with fewer confirming emission lines but a confident redshift ($z=0.8775$) based on the [OII] doublet feature. (d) Galaxy with a redshift ($z=0.9173$) based solely on the [OII] doublet. (e) Galactic star at $z=0$. (f) Quasar with a high-quality redshift ($z=0.3048$).

(Caltech). Contact: cblake@astro.swin.edu.au
We wish to acknowledge financial support for the WiggleZ project from The Australian Research Council, Swinburne University of Technology, The University of Queensland, the Anglo-Australian Observatory, and The Gregg Thompson Dark Energy Travel Fund.

References

- Bassett B *et al.* 2005 *A&G* **46** 5.26.
Bennett *et al.* 2003 *ApJS* **148** 1.
Blake C and Glazebrook K 2003 *ApJ* **594** 665.
Calder L and Lahav O 2008 *A&G* **49** 1.13.
Efstathiou G *et al.* 1990 *Nature* **348** 705.
Eisenstein D *et al.* 1999 *ApJ* **518** 2.
Eisenstein D *et al.* 2005 *ApJ* **633** 560.
Komatsu E *et al.* 2008 <http://arxiv.org/abs/0803.0547>.
Lewis I *et al.* 2002 *MNRAS* **333** 279.
Ostriker J and Steinhardt P 1995 *Nature* **377** 600.
Percival W *et al.* 2007 *MNRAS* **381** 1053.
Perlmutter S *et al.* 1999 *ApJ* **517** 565.
Riess A *et al.* 1998 *AJ* **116** 1009.
Saunders W *et al.* 2004 *SPIE* **5492** 389.
Seo H-J and Eisenstein D 2003 *ApJ* **598** 720.
Sharp R *et al.* 2006 *SPIE* **6269E** 14 <http://arxiv.org/abs/astro-ph/0606137>.
Spergel D *et al.* 2003 *ApJS* **148** 175.
Wiltshire D 2007 *Physics Rev. Lett.* **99** 1101.

FURTHER INFORMATION

2dFGRS

<http://www.mso.anu.edu.au/2dFGRS>

BOSS

<http://www.sdss3.org/cosmology.php>

Dark Energy Survey

<http://www.darkenergysurvey.org>

DEEP2

<http://deep.berkeley.edu>

GALEX

<http://www.galex.caltech.edu>

HETDEX

<http://www.as.utexas.edu/hetdex>

JDEM

<http://universe.nasa.gov/program/probes/jdem.html>

LSST

http://www.lsst.org/lstst_home.shtml

Panstarrs

<http://pan-starrs.ifa.hawaii.edu/public>

Planck

<http://www.rssd.esa.int/index.php?project=planck>

RCS2

<http://www.rcs2.org>

SDSS

<http://www.sdss.org>

Square Kilometre Array

http://www.skatelescope.org/pages/page_astronom.htm